

Dechant, Pierre-Philippe ORCID:

<https://orcid.org/0000-0002-4694-4010> (2014) What Clifford algebra can do for Coxeter groups and root systems. In: Maths HEP Lunchtime Seminar, 21st February 2014, Durham University. (Unpublished)

Downloaded from: <http://ray.yorks.ac.uk/id/eprint/4014/>

Research at York St John (RaY) is an institutional repository. It supports the principles of open access by making the research outputs of the University available in digital form. Copyright of the items stored in RaY reside with the authors and/or other copyright owners. Users may access full text items free of charge, and may download a copy for private study or non-commercial research. For further reuse terms, see licence terms governing individual outputs. [Institutional Repository Policy Statement](#)

# RaY

Research at the University of York St John

For more information please contact RaY at [ray@yorks.ac.uk](mailto:ray@yorks.ac.uk)



# What Clifford algebra can do for Coxeter groups and root systems

Pierre-Philippe Dechant

CPT, Mathematics Department, Durham University

Durham Mathematics HEP seminar – February 21, 2014

# The general theme: Geometry & Symmetry and their Applications

- Worked on a few different things: HEP – strings, particles and cosmology, pure maths and mathematical biology and Clifford algebras and mathematical physics
- Unifying themes of symmetry and geometry (euclidean, conformal, hyperbolic, spherical)
- Continuous Lie groups, e.g. for modeling cosmological spacetimes (Bianchi models)
- Discrete Coxeter groups and Kac-Moody algebras describe gravitational singularities/hidden symmetries in HEP theory, viruses, fullerenes, &c
- Mathematical frameworks of Coxeter groups and Clifford algebras

## 1 Introduction

- Coxeter groups and root systems
- Clifford algebras


## 2 Coxeter and Clifford

- The Induction Theorem – from 3D to 4D
- The Coxeter Plane
- Conformal Geometry
- Some Group Theory

## 3 Moonshine and Outlook



# Root systems – $A_2$




**Root system**  $\Phi$ : set of  
vectors  $\alpha$  such that

$$\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$$

and  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$

# Root systems – $A_2$



**Root system**  $\Phi$ : set of vectors  $\alpha$  such that

$$\Phi \cap \mathbb{R}\alpha = \{-\alpha, \alpha\} \quad \forall \alpha \in \Phi$$

and  $s_\alpha \Phi = \Phi \quad \forall \alpha \in \Phi$

**Simple roots**: express every element of  $\Phi$  via a  **$\mathbb{Z}$ -linear combination** (with coefficients of the same sign).

# Cartan Matrices

Cartan matrix of  $\alpha_i$ s is 
$$A_{ij} = 2 \frac{(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} = 2 \frac{|\alpha_j|}{|\alpha_i|} \cos \theta_{ij}$$

# Cartan Matrices

Cartan matrix of  $\alpha_i$ s is  $A_{ij} = 2 \frac{(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} = 2 \frac{|\alpha_j|}{|\alpha_i|} \cos \theta_{ij}$

$$\cos^2 \theta_{ij} = \frac{1}{4} A_{ij} A_{ji}$$

$$l_j^2 = \frac{A_{ij}}{A_{ji}} l_i^2$$

$$A_{ii} = 2$$

$$A_{ij} \in \mathbb{Z}^{\leq 0}$$

$$A_{ij} = 0 \Leftrightarrow A_{ji} = 0.$$

$$A_2: A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$$

# Cartan Matrices

Cartan matrix of  $\alpha_i$ s is  $A_{ij} = 2 \frac{(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} = 2 \frac{|\alpha_j|}{|\alpha_i|} \cos \theta_{ij}$

$$\cos^2 \theta_{ij} = \frac{1}{4} A_{ij} A_{ji}$$

$$l_j^2 = \frac{A_{ij}}{A_{ji}} l_i^2$$

$$A_{ii} = 2$$

$$A_{ij} \in \mathbb{Z}^{\leq 0}$$

$$A_{ij} = 0 \Leftrightarrow A_{ji} = 0.$$

$$A_2: A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$$

**Coxeter-Dynkin diagrams:** node = simple root, no link = roots orthogonal, simple link = roots at  $\frac{\pi}{3}$ , link with label  $m$  = angle  $\frac{\pi}{m}$ .

$$A_2 \circ \text{---} \circ$$

$$H_2 \circ \text{---}^5 \circ$$

$$I_2(n) \circ \text{---}^n \circ$$

# Coxeter groups

A **Coxeter group** is a group generated by some **involutive generators**  $s_i, s_j \in S$  subject to relations of the form  $(s_i s_j)^{m_{ij}} = 1$  with  $m_{ij} = m_{ji} \geq 2$  for  $i \neq j$ .

# Coxeter groups

A **Coxeter group** is a group generated by some **involutive generators**  $s_i, s_j \in S$  subject to relations of the form  $(s_i s_j)^{m_{ij}} = 1$  with  $m_{ij} = m_{ji} \geq 2$  for  $i \neq j$ .

The **finite** Coxeter groups have a **geometric representation** where the involutions are realised as **reflections** at **hyperplanes through the origin** in a Euclidean vector space  $\mathcal{E}$ . In particular, let  $(\cdot|\cdot)$  denote the inner product in  $\mathcal{E}$ , and  $v, \alpha \in \mathcal{E}$ .

The **generator**  $s_\alpha$  corresponds to the **reflection**

$$s_\alpha : v \rightarrow s_\alpha(v) = v - 2 \frac{(v|\alpha)}{(\alpha|\alpha)} \alpha$$

at a hyperplane perpendicular to the **root vector**  $\alpha$ .

The action of the **Coxeter group** is to permute these **root vectors**.

# Coxeter groups vs Lie groups vs Lie algebras vs root systems

- Lie group = **group** and **manifold**
- Lie algebra = **bilinear, antisymmetric bracket** and **Jacobi identity**
- Lie algebras are **infinitesimal** version of Lie group = near the identity
- Can be more comprehensive e.g. 2D conformal algebra vs 2D conformal group
- But finite group transformation laws can be easier than linearising
- 'Nice' Lie algebras have **triangular decomposition**:  
$$\mathcal{N}_- \oplus \mathcal{H} \oplus \mathcal{N}_+$$



# Coxeter groups vs Lie groups vs Lie algebras vs root systems

- 'Nice' Lie algebras have **triangular decomposition**:  
 $\mathcal{N}_- \oplus \mathcal{H} \oplus \mathcal{N}_+$
- $\mathcal{H}$  is the **Cartan subalgebra** (maximal commuting = quantum numbers)
- Creation and annihilation algebras  $\mathcal{N}$  form **root lattice**
- Symmetry group is called **Weyl group** and is a **crystallographic** Coxeter group:  $A_n, B_n/C_n, D_n, G_2, F_4, E_6, E_7, E_8$
- So Coxeter groups in theoretical physics always crystallographic! **Neglect**  $I_2(n), H_3, H_4$ .
- Useful Lie algebras are **(semi-)simple** LA (determinant of Cartan matrix is positive), **affine** LA (determinant is 0), **Kac-Moody algebras**, Borcherd's algebras...

# Kac-Moody algebras


- Kac-Moody algebras  $\mathcal{A}$  of rank  $N$  are defined by **generalised Cartan**  $(N \times N)$  matrices with  $A_{ii} = 2$ ,  $A_{ij} \in \mathbb{Z}_-(i \neq j)$  and  $A_{ij} \neq 0 \Rightarrow A_{ji} \neq 0$
- $3N$  **generators**  $h_i, e_i, f_i$  satisfy **Chevalley-Serre relations**  
 $[h_i, h_j] = 0$   $[h_i, e_j] = A_{ij}e_j$ ,  $[h_i, f_j] = -A_{ij}f_j$ ,  $[e_i, f_j] = \delta_{ij}h_i$   
 $\underbrace{[e_i, [e_i, [\dots, [e_i, e_j]]]] \dots]}_{1-A_{ij} \text{ times}} = 0$ ,  $\underbrace{[f_i, [f_i, [\dots, [f_i, f_j]]]] \dots]}_{1-A_{ij} \text{ times}} = 0$
- **Simple roots**  $\alpha_i$  are  $[h, e_i] = \alpha_i(h)e_i$

## Example – $A_1$ , $SU(2)$ , Angular Momentum



- Cartan subalgebra = Quantum number:  $L_z$
- $\mathcal{N}_+$ : raising operator  $L_+ = \alpha$
- $\mathcal{N}_-$ : lowering operator  $L_- = -\alpha$
- ( $L^2$  is Casimir/commutes with all algebra elements, is however not actually in the algebra!)

## Example – $A_1$ , $SU(2)$ , Electroweak



- Cartan subalgebra – Quantum number:  $A$
- $\mathcal{N}_+$ : raising operator  $W^+ = \alpha$
- $\mathcal{N}_-$ : lowering operator  $W^- = -\alpha$
- (Since SM electroweak is actually  $SU(2) \times U(1)$ ,  $U(1)$  gives another field  $i$ , such that physical  $Z^0$  and  $\gamma$  are superpositions of  $A$  and  $i$ )
- Also  $W^\pm$  now charged and self-interact, unlike QED

# Affine extensions


An **affine Coxeter group** is the extension of a Coxeter group by an **affine reflection in a hyperplane not containing the origin**  $s_{\alpha_0}^{aff}$  whose geometric action is given by

$$s_{\alpha_0}^{aff} v = \alpha_0 + v - \frac{2(\alpha_0 | v)}{(\alpha_0 | \alpha_0)} \alpha_0$$


**Non-distance preserving:** includes the **translation generator**

$$T v = v + \alpha_0 = s_{\alpha_0}^{aff} s_{\alpha_0} v$$

## Affine extensions – $A_2$




## Affine extensions – $A_2$





## Affine extensions – $A_2$

Affine extensions of crystallographic Coxeter groups lead to a **tessellation** of the plane and a **lattice**.






# Non-crystallographic Coxeter groups $H_2 \subset H_3 \subset H_4$



$$A = \begin{pmatrix} 2 & -\tau \\ -\tau & 2 \end{pmatrix}$$



$$A = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -\tau \\ 0 & -\tau & 2 \end{pmatrix}$$



$$A = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -\tau \\ 0 & 0 & -\tau & 2 \end{pmatrix}$$

$H_2 \subset H_3 \subset H_4$ : 10, 120, 14,400 elements, the only Coxeter groups that generate **rotational symmetries of order 5**  
linear combinations now in the **extended integer ring**

$$\mathbb{Z}[\tau] = \{a + \tau b \mid a, b \in \mathbb{Z}\} \quad \text{golden ratio}$$

$$\tau = \frac{1}{2}(1 + \sqrt{5}) = 2 \cos \frac{\pi}{5}$$

$$x^2 = x + 1$$

$$\tau' = \sigma = \frac{1}{2}(1 - \sqrt{5}) = 2 \cos \frac{2\pi}{5}$$


$$\tau + \sigma = 1, \tau\sigma = -1$$

# What's new?

- In HEP, mostly come from Lie groups, then Lie algebras, then their Weyl groups and root systems
- This only gives the crystallographic Coxeter groups
- Do the non-crystallographic Coxeter groups have something interesting to offer? In particular, affine extensions?
- Interesting connections between the geometries of different dimensions: Relation between crystallographic and non-crystallographic ( $E_8$  and  $H_4$ ) and my spinor construction (3 & 4D)
- Both could have interesting consequences for HEP (4D groups and  $E_8$  feature heavily) and other applications (viruses, quasicrystals, proteins, fullerenes...)



# Affine extensions of non-crystallographic root systems

Unit translation along a vertex of a unit pentagon




# Affine extensions of non-crystallographic root systems

Unit translation along a vertex of a unit pentagon



# Affine extensions of non-crystallographic root systems


Unit translation along a vertex of a unit pentagon



A **random** translation would give 5 secondary pentagons, i.e. 25 points. Here we have **degeneracies** due to 'coinciding points'.


# Affine extensions of non-crystallographic root systems

Translation of length  $\tau = \frac{1}{2}(1 + \sqrt{5}) \approx 1.618$  (golden ratio)




Looks like a **virus** or **carbon onion**

# More Blueprints




# Road Map





# Applications of affine extensions of non-crystallographic root systems



There are interesting applications to **quasicrystals**, **viruses** or **carbon onions**, but here concentrate on the **mathematical** aspects

# Basics of Clifford Algebra I

- Form an algebra using the **Geometric Product**

$$ab \equiv a \cdot b + a \wedge b \text{ for two vectors}$$

- Extend via linearity and associativity to higher grade elements (**multivectors**)
- For an n-dimensional space generated by n orthogonal unit vectors  $e_i$  have  $2^n$  elements
- Then  $e_i e_j = e_i \wedge e_j = -e_j e_i$  so **anticommute** (Grassmann variables, exterior algebra)
- Unlike the inner and outer products separately, this product is **invertible**
- This feeds through to the **differential structure** of the theory with more powerful **Greens functions methods**  $\nabla^{-1}$

## Basics of Clifford Algebra II

- These are known to have **matrix representations** over the normed division algebras  $\mathbb{R}$ ,  $\mathbb{C}$  and  $\mathbb{H} \Rightarrow$  **Classification** of Clifford algebras
- E.g. **Pauli algebra** in 3D (likewise for **Dirac algebra** in 4D) is

$$\underbrace{\{1\}}_{1 \text{ scalar}} \quad \underbrace{\{e_1, e_2, e_3\}}_{3 \text{ vectors}} \quad \underbrace{\{e_1 e_2, e_2 e_3, e_3 e_1\}}_{3 \text{ bivectors}} \quad \underbrace{\{I \equiv e_1 e_2 e_3\}}_{1 \text{ trivector}}$$

- These have the well-known matrix representations in terms of  **$\sigma$ - and  $\gamma$ -matrices**
- Working with these is not necessarily the most insightful thing to do, so here stress approach to **work directly** with the algebra
- Naturally have things that **square to  $-1$** , e.g.

$$\boxed{(e_1 e_2)^2 = e_1 e_2 e_1 e_2 = -e_1 e_1 e_2 e_2 = -1}, \text{ and } \text{non-trivial commutation properties}$$

# Reflections

- Clifford algebra is **very efficient** at performing **reflections**
- Consider reflecting the vector  $a$  in a hypersurface with unit normal  $n$ :

$$a' = a_{\perp} - a_{\parallel} = a - 2a_{\parallel} = a - 2(a \cdot n)n$$

- c.f. **fundamental Weyl reflection**  $s_i : v \rightarrow s_i(v) = v - 2 \frac{(v|\alpha_i)}{(\alpha_i|\alpha_i)} \alpha_i$
- But in Clifford algebra have  $n \cdot a = \frac{1}{2}(na + an)$  so reassembles into **sandwiching**

$$a' = -nan$$

- So both **Coxeter** and **Clifford** frameworks are ideally suited to describing **reflections** – first to combine the two

# Reflections and Rotations

- Generate a **rotation** when compounding two reflections wrt  $n$  then  $m$  (**Cartan-Dieudonné theorem**):

$$a'' = mn an m \equiv R a \tilde{R}$$

where  $R = mn$  is called a **rotor** and a tilde denotes **reversal** of the order of the constituent vectors ( $R \tilde{R} = 1$ )

- Now neat thing is all multivectors transform **covariantly** e.g.

$$MN \rightarrow (R M \tilde{R})(R N \tilde{R}) = R M \tilde{R} R N \tilde{R} = R(MN) \tilde{R}$$

so transform **double-sidedly**

- Rotors form a **group**, the rotor group, which gives a representation of the **Spin group**  $Spin(n)$  – they transform **single-sidedly** (obvious now it's a double (universal) cover)

# Artin's Theorem and orthogonal transformations

- **Artin**: every isometry is at most  $d$  reflections
- Since have a **double cover** of reflections ( $n$  and  $-n$ ) we have a **double cover** of  $O(p, q)$ :  **$\text{Pin}(p, q)$**

$$x' = \pm n_1 n_2 \dots n_k x n_k \dots n_2 n_1$$

- Pinors/vectors = products of vectors  $n_1 n_2 \dots n_k$  encode orthogonal transformations via '**sandwiching**'
- **Cartan-Dieudonné**: rotations are an even number of reflections:  **$\text{Spin}(p, q)$**  doubly covers  $SO(p, q)$
- The conformal group  **$C(p, q) \sim SO(p+1, q+1)$**  so can use these for **translations, inversions** etc as well

# Spinor techniques

- Of course there is a **matrix representation**  $\underline{R}$  for the action of a **spinor**:  $\underline{R}x = Rx\tilde{R}$
- This is the usual **rotation matrix**  $\underline{R}$  in  $SO(p, q)$
- Having the **spin double cover/square root of the rotation matrix** can be convenient
- E.g. can get **differential equations for spinor**  $R$  that are **easier** to solve, then can reconstitute  $\underline{R}$  if necessary

## 1 Introduction

- Coxeter groups and root systems
- Clifford algebras

## 2 Coxeter and Clifford

- The Induction Theorem – from 3D to 4D
- The Coxeter Plane
- Conformal Geometry
- Some Group Theory

## 3 Moonshine and Outlook



## 3D Platonic Solids



- There are 5 Platonic solids
- Tetrahedron (**self-dual**) ( $A_3$ )
- **Dual** pair **octahedron** and **cube** ( $B_3$ )
- **Dual** pair **icosahedron** and **dodecahedron** ( $H_3$ )
- Only the **octahedron** is a **root system** (actually for  $(A_1^3)$ )

# Clifford and Coxeter: Platonic Solids



Platonic Solid	Group	root system
Tetrahedron	$A_3$ $A_1^3$	Cuboctahedron Octahedron
Octahedron Cube	$B_3$	Cuboctahedron + Octahedron
Icosahedron Dodecahedron	$H_3$	Icosidodecahedron

- **Platonic Solids** have been known for millennia

# Clifford and Coxeter: Platonic Solids



$A_3$

$A_1$

$B_3$

$H_3$

Platonic Solid	Group	root system
Tetrahedron	$A_3$ $A_1^3$	Cuboctahedron Octahedron
Octahedron Cube	$B_3$	Cuboctahedron + Octahedron
Icosahedron Dodecahedron	$H_3$	Icosidodecahedron

- **Platonic Solids** have been known for millennia
- Described by **Coxeter** groups


# Clifford and Coxeter: Platonic Solids



$A_1^3$	$A_1^4$
$A_3$	$D_4$
$B_3$	$F_4$
$H_3$	$H_4$

- **Platonic Solids** have been known for millennia; described by **Coxeter** groups
- Concatenating reflections gives **Clifford** spinors (**binary polyhedral groups**)
- These **induce 4D root systems**  

$$\psi = a_0 + a_i |e_i| \Rightarrow \psi \tilde{\psi} = a_0^2 + a_1^2 + a_2^2 + a_3^2$$
- 4D analogues of the Platonic Solids and give rise to 4D **Coxeter** groups



## 4D 'Platonic Solids'

- In 4D, there are **6 analogues** of the Platonic Solids:
- **5-cell** (self-dual) ( $A_4$ )
- **24-cell** (self-dual) ( $D_4$ ) – a 24-cell and its dual together are the  **$F_4$  root system**
- Dual pair **16-cell** and **8-cell** ( $B_4$ )
- Dual pair **600-cell** and **120-cell** ( $H_4$ )
- **24-cell, 16-cell and 600-cell** are all **root systems**, as is the related  **$F_4$  root system**
- 8-cell and 120-cell are dual to a root system, so in 4D **out of 6 Platonic Solids only the 5-cell** (corresponding to  $A_n$  family) is not related to a root system!
- The 4D Platonic solids are **not normally thought to be related to the 3D ones** except for the boundary cells

# Spinorial Symmetries of 4D Polytopes

## Spinorial symmetries

rank 3	$ \Phi $	$ W $	rank 4	$ \Phi $	Symmetry
$A_3$	12	24	$D_4$ 24-cell	24	$2 \cdot 24^2 = 576$
$B_3$	18	48	$F_4$ lattice	48	$48^2 = 2304$
$H_3$	30	120	$H_4$ 600-cell	120	$120^2 = 14400$
$A_1^3$	6	8	$A_1^4$ 16-cell	8	$3! \cdot 8^2 = 384$
$A_1 \oplus A_2$	8	12	$A_2 \oplus A_2$ prism	12	$12^2 = 144$
$A_1 \oplus H_2$	12	20	$H_2 \oplus H_2$ prism	20	$20^2 = 400$
$A_1 \oplus I_2(n)$	$n+2$	$2n$	$I_2(n) \oplus I_2(n)$	$2n$	$(2n)^2$

Similar for **Grand Antiprism** ( $H_4$  without  $H_2 \oplus H_2$ ) and **Snub 24-cell** ( $2I$  without  $2T$ ).

# Induction Theorem

- Theorem: 3D spinor groups are root systems ( $R$  and  $-R$  are in a spinor group by construction, and closure under reflections is guaranteed by the closure property of the spinor group)
- Induction Theorem: Every rank-3 root system induces a rank-4 root system.
- Counterexample: not every rank-4 root system is induced in this way
- Spinor group is trivially closed under conjugation, left and right multiplication. Results in non-trivial symmetries when viewed as a polytope/root system.
- Now explains symmetry of the polytopes/root system and thus the order of the rank-4 Coxeter group

# Induction Theorem

- So induced **4D polytopes** are actually **root systems**.
- Clear why the **number of roots**  $|\Phi|$  is equal to  $|G|$ , the **order of the spinor group**
- Theorem: The **automorphism group** of the induced root system contains **two factors** of the respective spinor group acting from the **left** and the **right**.
- Only **remaining cases** in 3D are  $A_1 \oplus I_2(n)$ , which give  $I_2(n) \oplus I_2(n)$



# General Case of Induction

Only **remaining case** is what happens for  $A_1 \oplus I_2(n)$  - this gives a **doubling**  $I_2(n) \oplus I_2(n)$

rank 3	rank 4
$A_3$	$D_4$
$B_3$	$F_4$
$H_3$	$H_4$
$A_1^3$	$A_1^4$
$A_1 \oplus A_2$	$A_2 \oplus A_2$
$A_1 \oplus H_2$	$H_2 \oplus H_2$
$A_1 \oplus I_2(n)$	$I_2(n) \oplus I_2(n)$

# Spinorial Symmetries of 4D Polytopes

## Spinorial symmetries

rank 3	$ \Phi $	$ W $	rank 4	$ \Phi $	Symmetry
$A_3$	12	24	$D_4$ 24-cell	24	$2 \cdot 24^2 = 576$
$B_3$	18	48	$F_4$ lattice	48	$48^2 = 2304$
$H_3$	30	120	$H_4$ 600-cell	120	$120^2 = 14400$
$A_1^3$	6	8	$A_1^4$ 16-cell	8	$3! \cdot 8^2 = 384$
$A_1 \oplus A_2$	8	12	$A_2 \oplus A_2$ prism	12	$12^2 = 144$
$A_1 \oplus H_2$	12	20	$H_2 \oplus H_2$ prism	20	$20^2 = 400$
$A_1 \oplus I_2(n)$	$n+2$	$2n$	$I_2(n) \oplus I_2(n)$	$2n$	$(2n)^2$

Similar for **Grand Antiprism** ( $H_4$  without  $H_2 \oplus H_2$ ) and **Snub 24-cell** ( $2I$  without  $2T$ ). Additional factors in the automorphism group come from **3D Dynkin diagram symmetries**!

## Some non-Platonic examples of spinorial symmetries

- **Grand Antiprism**: the 100 vertices achieved by subtracting 20 vertices of  $H_2 \oplus H_2$  from the 120 vertices of the  $H_4$  root system 600-cell – two separate orbits of  $H_2 \oplus H_2$
- This is a semi-regular polytope with automorphism symmetry  $\text{Aut}(H_2 \oplus H_2)$  of order  $400 = 20^2$
- Think of the  $H_2 \oplus H_2$  as coming from the **doubling procedure**? (Likewise for  $\text{Aut}(A_2 \oplus A_2)$  subgroup)
- **Snub 24-cell**:  $2T$  is a subgroup of  $2I$  so subtracting the 24 corresponding vertices of the 24-cell from the 600-cell, one gets a semiregular polytope with 96 vertices and automorphism group  $2T \times 2T$  of order  $576 = 24^2$ .

# Sub root systems

- The above spinor groups had spinor multiplication as the **group operation**
- But also closed under **twisted conjugation** – corresponds to **closure under reflections** (root system property)
- If we take **twisted conjugation** as the group operation instead, we can have various **subgroups**
- These are the remaining **4D root systems** e.g.  $A_4$  or  $B_4$

# What's new?

- Novel **connection** between geometry of **3D and 4D**
- In fact, 3D seems more **fundamental** – contrary to the **usual perspective** of 3D subgroups of 4D groups
- **Spinorial symmetries**
- Clear why **spinor group** gives a root system and why **two factors** of the same group reappear in the **automorphism group**
- Novel **spinorial perspective** on 4D geometry
- **Accidentalness** of the spinor construction and **exceptional** 4D phenomena
- Connection with Arnold's **trinities**, the **McKay correspondence** and **Monstrous Moonshine**

## Recap: Clifford algebra and reflections & rotations

- Clifford algebra is **very efficient** at performing **reflections** via **sandwiching**

$$a' = -nan$$




- Generate a **rotation** when compounding two reflections wrt  $n$  then  $m$  (**Cartan-Dieudonné theorem**):

$$a'' = mn an m \equiv R a \tilde{R}$$

where  $R = mn$  is called a **rotor** and a tilde denotes **reversal** of the order of the constituent vectors ( $R\tilde{R} = 1$ )

## From the Coxeter simple roots to the root system

- Take the  $A_1 \times A_1 \times A_1$  simple roots  $(1,0,0)$ ,  $(0,1,0)$ ,  $(0,0,1)$   
 $\Rightarrow$  under reflections get  $(-1,0,0)$ ,  $(0,-1,0)$ ,  $(0,0,-1)$ , the vertices of an **octahedron**.
- Take the **three simple roots** of  $A_1 \times A_1 \times A_1 / A_3 / B_3 / H_3$ .  
Closure under Clifford **reflections** generate the whole root system of 6/12/18/30 **vertices of an octahedron/cuboctahedron/ cuboctahedron with an octahedron/ icosidodecahedron**).



## Spinors from reflections

- These are the 3D Coxeter groups that are symmetry groups of the **Platonic Solids** (tetrahedron and octahedron are similar but simpler than the icosahedron)
- The 6/12/18/30 **reflections** in  $A_1 \times A_1 \times A_1 / A_3 / B_3 / H_3$  generate 8/24/48/120 **rotors**.
- E.g.  $(\pm 1, 0, 0)$ ,  $(0, \pm 1, 0)$ ,  $(0, 0, \pm 1)$  give the 8 permutations of  $(\pm 1; 0, 0, 0)$  (scalar and bivector parts, the notation will become clear later).
- The **discrete spinor group** is isomorphic to the **quaternion group**  $Q$  / **binary tetrahedral group**  $2T$  / **binary octahedral group**  $2O$  / **binary icosahedral group**  $2I$ ).







# A unified framework for polyhedral groups

Group	Discrete subgroup	Action Mechanism
$SO(3)$	rotational (chiral)	$x \rightarrow \tilde{R}xR$
$O(3)$	reflection (full/Coxeter)	$x \rightarrow \pm \tilde{A}xA$
$Spin(3)$	binary	$(R_1, R_2) \rightarrow R_1 R_2$
$Pin(3)$	pinor	$(A_1, A_2) \rightarrow A_1 A_2$

- e.g. the **chiral icosahedral** group has 60 elements, encoded in Clifford by 120 rotors, which form the **binary icosahedral** group
- together with the **inversion/pseudoscalar**  $I$  this gives 60 rotations and 60 rotoinversions, i.e. the **full icosahedral** group  $H_3$  in 120 elements (with 240 versors)
- all three are interesting groups, e.g. in **neutrino and flavour physics** for family symmetry model building

# Spinors and Polytopes

- The space of  $Cl(3)$ -spinors and quaternions have a **4D Euclidean signature**:  $\psi = a_0 + a_i |e_i \Rightarrow \psi \tilde{\psi} = a_0^2 + a_1^2 + a_2^2 + a_3^2$
- Can reinterpret **spinors in  $\mathbb{R}^3$**  as **vectors in  $\mathbb{R}^4$**
- Then the spinors constitute the **vertices** of the **16-cell**, **24-cell**, **24-cell and dual 24-cell** and the **600-cell**
- These are 4D analogues of the **Platonic Solids**: regular convex 4-polytopes



# Spinors, Polytopes and Root systems

- The 16-cell, 24-cell, 24-cell and dual 24-cell and the 600-cell are in fact the root systems of  $A_1 \times A_1 \times A_1 \times A_1$ ,  $D_4$ ,  $F_4$  and  $H_4$
- Exceptional phenomena:  $D_4$  (triality, important in string theory),  $F_4$  (largest lattice symmetry in 4D),  $H_4$  (largest non-crystallographic symmetry)
- Exceptional  $D_4$  and  $F_4$  arise from series  $A_3$  and  $B_3$
- In fact, can strengthen this statement on inducing polytopes to statement on inducing root systems

# Root systems in three and four dimensions

The **spinors** generated from the reflections contained in the respective **rank-3 Coxeter group** via the geometric product are realisations of the **binary polyhedral groups**  $Q$ ,  $2T$ ,  $2O$  and  $2I$ , which were known to generate (mostly exceptional) **rank-4 groups**, but **not known why**, and why the ‘**mysterious symmetries**’.

rank-3 group	diagram	binary	rank-4 group	diagram
$A_1 \times A_1 \times A_1$	○ ○ ○	$Q$	$A_1 \times A_1 \times A_1 \times A_1$	○ ○ ○ ○
$A_3$	○—○—○	$2T$	$D_4$	○—○—○   ○
$B_3$	○—○—○ <sup>4</sup>	$2O$	$F_4$	○—○—○—○ <sup>4</sup>
$H_3$	○—○—○ <sup>5</sup>	$2I$	$H_4$	○—○—○—○ <sup>5</sup>

# Arnold's Trinities

Arnold's observation that many areas of real mathematics can be **complexified** and **quaternionified** resulting in theories with a similar structure.

- The **fundamental trinity** is thus  $(\mathbb{R}, \mathbb{C}, \mathbb{H})$
- The **projective spaces**  $(\mathbb{R}P^n, \mathbb{C}P^n, \mathbb{H}P^n)$
- The **spheres**  $(\mathbb{R}P^1 = S^1, \mathbb{C}P^2 = S^2, \mathbb{H}P^1 = S^4)$
- The **Möbius/Hopf bundles**  $(S^1 \rightarrow S^1, S^4 \rightarrow S^2, S^7 \rightarrow S^4)$
- The **Lie Algebras**  $(E_6, E_7, E_8)$
- The symmetries of the **Platonic Solids**  $(A_3, B_3, H_3)$
- The **4D groups**  $(D_4, F_4, H_4)$
- **New connections** via my **Clifford spinor construction** (see McKay correspondence)


# Platonic Trinities

- Arnold's connection between  $(A_3, B_3, H_3)$  and  $(D_4, F_4, H_4)$  is **very convoluted** and involves numerous other trinities at intermediate steps:
- **Decomposition of the projective plane** into Weyl chambers and Springer cones
- The **number of Weyl chambers** in each segment is  $24 = 2(1 + 3 + 3 + 5)$ ,  $48 = 2(1 + 5 + 7 + 11)$ ,  $120 = 2(1 + 11 + 19 + 29)$
- Notice this miraculously **matches the quasihomogeneous weights**  $((2, 4, 4, 6), (2, 6, 8, 12), (2, 12, 20, 30))$  of the Coxeter groups  $(D_4, F_4, H_4)$
- Believe the Clifford connection is **more direct**

## Some Group Theory: chiral, full, binary, pin

- Easy enough to calculate **conjugacy classes** etc of versors in Clifford
- Chiral (**binary**) polyhedral groups have irreps
- tetrahedral (12/24):  $1, 1', 1'', 2_s, 2'_s, 2''_s, 3$
- octahedral (24/48):  $1, 1', 2, 2_s, 2'_s, 3, 3', 4_s$
- icosahedral (60/120):  $1, 2_s, 2'_s, 3, \bar{3}, 4, 4_s, 5, 6_s$
- All binary are **discrete subgroups of  $SU(2)$**  and all thus have a  $2_s$  spinor irrep
- Connection with the **McKay correspondence!**

# Affine extensions – $E_8^-$



$$-\alpha_0 = 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 5\alpha_4 + 6\alpha_5 + 4\alpha_6 + 2\alpha_7 + 3\alpha_8$$

AKA  $E_8^+$  and along with  $E_8^{++}$  and  $E_8^{+++}$  thought to be the underlying symmetry of **String and M-theory**

Also interesting from a pure mathematics point of view:  **$E_8$  lattice**, **McKay correspondence** and **Monstrous Moonshine**.




# The McKay Correspondence


binary polyhe-  
dral groups  
 $2T, 2O, 2I$   
 $\sum d_i$  12, 18, 30  
 $\sum d_i^2$  24, 48, 120

McKay correspondence

Exceptional  
Lie Groups  
 $E_6$ , 12  
 $E_7$ , 18  
 $E_8$ , 30  
(Coxeter numbers)



# The McKay Correspondence



# The McKay Correspondence

More than E-type groups: the infinite family of 2D groups, the **cyclic** and **dicyclic groups** are in correspondence with  $A_n$  and  $D_n$ , e.g. the quaternion group  $Q$  and  $D_4^+$ . So McKay correspondence not just a trinity but **ADE-classification**. We also have  $I_2(n)$  on top of the trinity ( $A_3, B_3, H_3$ )

rank-3 group	diagram	binary	rank-4 group	diagram	Lie algebra	diagram
$A_1 \times A_1 \times A_1$		$Q$	$A_1 \times A_1 \times A_1 \times A_1$		$D_4^+$	
$A_3$		$2T$	$D_4$		$E_6^+$	
$B_3$		$2O$	$F_4$		$E_7^+$	
$H_3$		$2I$	$H_4$		$E_8^+$	

## 4D geometry is surprisingly important for HEP

- 4D root systems are **surprisingly relevant to HEP**
- $A_4$  is  $SU(5)$  and comes up in **Grand Unification**
- $D_4$  is  $SO(8)$  and is the little group of **String theory**
- In particular, its **triality symmetry** is crucial for showing the equivalence of RNS and GS strings
- $B_4$  is  $SO(9)$  and is the little group of **M-Theory**
- $F_4$  is the **largest crystallographic** symmetry in 4D and  $H_4$  is the **largest non-crystallographic** group
- The above are **subgroups** of the latter two
- **Spinorial nature** of the root systems could have **surprising consequences for HEP**

# Quaternions and Clifford Algebra

- The unit **spinors**  $\{1; i; j; k\}$  of  $\text{Cl}(3)$  are isomorphic to the **quaternion** algebra  $\mathbb{H}$  (up to sign)
- The 3D **Hodge dual of a vector** is a **pure bivector** which corresponds to a **pure quaternion**, and their products are identical (up to sign)









# Discrete Quaternion groups

- The 8 quaternions of the form  $(\pm 1, 0, 0, 0)$  and permutations are called the **Lipschitz units**, and form a realisation of the **quaternion group** in 8 elements.
- The 8 Lipschitz units together with  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  are called the **Hurwitz units**, and realise the **binary tetrahedral group** of order 24. Together with the 24 'dual' quaternions of the form  $\frac{1}{\sqrt{2}}(\pm 1, \pm 1, 0, 0)$ , they form a group isomorphic to the **binary octahedral group** of order 48.
- The 24 Hurwitz units together with the 96 unit quaternions of the form  $(0, \pm \tau, \pm 1, \pm \sigma)$  and even permutations, are called the **Icosians**. The icosian group is isomorphic to the **binary icosahedral group** with 120 elements.

# Quaternionic representations of 3D and 4D Coxeter groups

- Groups  $E_8$ ,  $D_4$ ,  $F_4$  and  $H_4$  have representations in terms of **quaternions**
- **Extensively used** in the high energy physics/quasicrystal/Coxeter/polytope literature and thought of as deeply significant, though not really clear why
- e.g.  $H_4$  consists of 120 elements of the form  $(\pm 1, 0, 0, 0)$ ,  $\frac{1}{2}(\pm 1, \pm 1, \pm 1, \pm 1)$  and  $(0, \pm \tau, \pm 1, \pm \sigma)$
- Seen as remarkable that the **subset of the 30 pure quaternions** is a realisation of  $H_3$  (**a sub-root system**)
- Similarly,  $A_3$ ,  $B_3$ ,  $A_1 \times A_1 \times A_1$  have representations in terms of **pure quaternions**
- Will see there is a **much simpler geometric explanation**

# Quaternionic representations used in the literature





# Demystifying Quaternionic Representations

- 3D: **Pure quaternions** = Hodge dualised (pseudoscalar) **root vectors**
- In fact, they are the **simple roots of the Coxeter groups**
- 4D: **Quaternions** = disguised **spinors** – but those of the **3D Coxeter group** i.e. the binary polyhedral groups!
- This relation between 3D and 4D via the geometric product does not seem to be known
- Quaternion multiplication = ordinary Clifford reflections and rotations

# Demystifying Quaternionic Representations

- Pure quaternion subset of 4D groups only gives 3D group if the 3D group contains the inversion/pseudoscalar /
- e.g. does not work for the tetrahedral group  $A_3$ , but  $A_3 \rightarrow D_4$  induction still works, with the central node essentially 'spinorial'
- In fact, it goes the other way around: the 3D groups induce the 4D groups via spinors
- The rank-4 groups are also generated (under quaternion multiplication) by two quaternions we can identify as  $R_1 = \alpha_1 \alpha_2$  and  $R_2 = \alpha_2 \alpha_3$
- Can see these are 'spinor generators' and how they don't really contain any more information/roots than the rank-3 groups alone

## Quaternions vs Clifford versors

- **Sandwiching** is often seen as particularly nice feature of the **quaternions giving rotations**
- This is actually a **general feature** of Clifford algebras/versors **in any dimension**; the isomorphism to the **quaternions** is **accidental** to 3D
- However, the **root system** construction does not necessarily generalise
- 2D generalisation merely gives that  $I_2(n)$  is **self-dual**
- **Octonionic** generalisation just induces two copies of the above 4D root systems, e.g.  $A_3 \rightarrow D_4 \oplus D_4$

## References (single-author)

- Clifford algebra unveils a surprising geometric significance of quaternionic root systems of Coxeter groups  
Advances in Applied Clifford Algebras, June 2013, Volume 23, Issue 2, pp 301-321
- A Clifford algebraic framework for Coxeter group theoretic computations (Conference Prize at AGACSE 2012)  
Advances in Applied Clifford Algebras (2013)
- Nomination for W.K. Clifford Prize (2014)
- Invitation to Arizona State University
- Rank-3 root systems induce root systems of rank 4 via a new Clifford spinor construction arXiv:1207.7339 (2012)
- Platonic Solids generate their 4-dimensional analogues  
Acta Cryst. A69 (2013)

# Coxeter Elements, Degrees and Exponents

- Like the symmetric group, Coxeter groups can have **invariant polynomials**. Their **degrees**  $d$  are important invariants/group characteristics.
- Turns out that actually **degrees**  $d$  are intimately related to so-called **exponents**  $m$   $m = d - 1$ .

# Coxeter Elements, Degrees and Exponents




- A **Coxeter Element** is any combination of all the simple reflections  $w = s_1 \dots s_n$ , i.e. in Clifford algebra it is encoded by the versor  $W = \alpha_1 \dots \alpha_n$  acting as  $v \rightarrow wv = \pm \tilde{W} v W$ . All such elements are conjugate and thus their **order** is invariant and called the **Coxeter number  $h$** .
- The Coxeter element has **complex eigenvalues** of the form  $\exp(2\pi mi/h)$  where  $m$  are called **exponents**.
- Standard theory **complexifies** the real Coxeter group situation in order to find **complex eigenvalues**, then takes **real sections** again (the unfortunate standard procedure in many situations) – without any insight into the complex structure (or in fact, there are different ones).

# Coxeter Elements, Degrees and Exponents

- The Coxeter element has **complex eigenvalues** of the form  $\exp(2\pi mi/h)$  where  $m$  are called **exponents**
- Standard theory **complexifies** the real Coxeter group situation in order to find **complex eigenvalues**, then takes **real** sections again (the unfortunate standard procedure in many situations) – without any insight into the complex structure(s)
- In particular, **1** and  **$h-1$**  are always exponents
- Turns out that actually **exponents and degrees** are intimately related ( $m = d - 1$ ). The construction is slightly roundabout but uniform, and uses the **Coxeter plane**.

# The Coxeter Plane


- Can show **every** (for our purposes) Coxeter group has a Coxeter plane.
- A way to visualise Coxeter groups in any dimension by **projecting** their root system onto the Coxeter plane





# The Coxeter Plane

- Obvious from Clifford point of view, that Coxeter element has eigenspaces (**eigenblades**) rather than just eigenvectors
- In particular, can show **every** (for our purposes) Coxeter group has a Coxeter plane
- Existence relies on the fact that all groups in question have **tree-like Dynkin diagrams**, and thus admit an **alternate colouring**
- Essentially just gives **two sets of mutually commuting generators**



# The Coxeter Plane

- Existence relies on the fact that all groups in question have **tree-like Dynkin diagrams**, and thus admit an alternate colouring
- Essentially just gives **two sets of orthogonal = mutually commuting generators but anticommuting root vectors**  $\alpha_w$  and  $\alpha_b$  (duals  $\omega$ )
- Cartan matrices are positive definite, and thus have a **Perron-Frobenius** (all positive) eigenvector  $\lambda_i$ .
- Take **linear combinations** of components of this eigenvector as coefficients of two vectors from the orthogonal sets  
$$v_w = \sum \lambda_w \omega_w \text{ and } v_b = \sum \lambda_b \omega_b$$
- Their **outer product/Coxeter plane bivector**  $B_C = v_b \wedge v_w$  describes an **invariant plane** where  $w$  acts by rotation by  $2\pi/h$ .

# Clifford Algebra and the Coxeter Plane – 2D case

- For  $I_2(n)$  take  $\alpha_1 = e_1, \alpha_2 = -\cos \frac{\pi}{n} e_1 + \sin \frac{\pi}{n} e_2$

- So Coxeter versor is just

$$W = \alpha_1 \alpha_2 = -\cos \frac{\pi}{n} + \sin \frac{\pi}{n} e_1 e_2 = -\cos \frac{\pi}{n} + \sin \frac{\pi}{n} I = -\exp \left( -\frac{\pi I}{n} \right)$$

- In Clifford algebra it is therefore immediately obvious that the action of the  $I_2(n)$  Coxeter element is described by a versor (here a rotor/spinor) that encodes rotations in the  $e_1 e_2$ -Coxeter-plane and yields  $h = n$  since trivially  $W^n = (-1)^{n+1}$  yielding  $w^n = 1$  via  $wv = \tilde{W}vW$ .

# Clifford Algebra and the Coxeter Plane – 2D case

- So **Coxeter versor** is just  $W = -\exp\left(-\frac{\pi I}{n}\right)$
- $I = e_1 e_2$  **anticommutes** with both  $e_1$  and  $e_2$  such that **sandwiching formula** becomes

$$v \rightarrow wv = \tilde{W}vW = \tilde{W}^2v = \exp\left(\pm \frac{2\pi I}{n}\right)v \text{ immediately}$$

yielding the standard result for the **complex eigenvalues** in real Clifford algebra **without any need for artificial complexification**

- The Coxeter plane bivector  $B_C = e_1 e_2 = I$  gives the **complex structure**
- The Coxeter plane bivector  $B_C$  is invariant under the **Coxeter versor**  $\tilde{W}B_CW = \pm B_C$ .

## Clifford Algebra and the Coxeter Plane – 3D case

- In 3D,  $A_3$ ,  $B_3$ ,  $H_3$  have  $\{1, 2, 3\}$ ,  $\{1, 3, 5\}$  and  $\{1, 5, 9\}$
- Coxeter element is product of a **spinor** in the Coxeter plane with the same complex structure as before, and a **reflection perpendicular** to the plane
- So in 3D still completely determined by the plane
- $1$  and  $h-1$  are **rotations** in **Coxeter plane**
- $h/2$  is the **reflection** (for  $v$  in the normal direction)

$$wv = \tilde{W}^2 = \exp\left(\pm \frac{2\pi I}{h} \frac{h}{2}\right) = \exp(\pm \pi I)v = -v$$

# Clifford algebra: no need for complexification

- Turns out in Clifford algebra we can **factorise**  $W$  into **orthogonal** (commuting/anticommuting) components  

$$W = \alpha_1 \dots \alpha_n = W_1 \dots W_n \text{ with } W_i = \exp(\pi m_i l_i / h)$$
- Here,  $l_i$  is a bivector describing a **plane** with  $l_i^2 = -1$
- For  $v$  **orthogonal to the plane** described by  $l_i$  we have  

$$v \rightarrow \tilde{W}_i v W_i = \tilde{W}_i W_i v = v \text{ so cancels out}$$
- For  $v$  **in the plane** we have  

$$v \rightarrow \tilde{W}_i v W_i = \tilde{W}_i^2 v = \exp(2\pi m_i l_i / h) v$$
- Thus if we **decompose**  $W$  into **orthogonal eigenspaces**, in the eigenvector equation all orthogonal bits cancel out and one gets the complex eigenvalue from the respective eigenspace

## Clifford algebra: no need for complexification

- For  $v$  in the plane we have


$$v \rightarrow \tilde{W}_i v W_i = \tilde{W}_i^2 v = \exp(2\pi m_i l_i / h) v$$

- So **complex eigenvalue equation** arises geometrically **without any need** for complexification
- **Different complex structures** immediately give different **eigenplanes**
- Eigenvalues/angles/**exponents** given from just factorising  $W = \alpha_1 \dots \alpha_n$
- E.g.  $B_4$  has exponents 1, 3, 5, 7 and  $W = \exp\left(\frac{\pi}{8} l_1\right) \exp\left(\frac{3\pi}{8} l_2\right)$
- Here we have been looking for orthogonal eigenspaces, so **innocuous** – different complex structures commute
- But not in general – **naive complexification** can be misleading

# Clifford Algebra and the Coxeter Plane – 4D case

- E.g.  $B_4$  has exponents 1, 3, 5, 7
- Coxeter versor decomposes into **orthogonal components**

$$W = \alpha_1 \alpha_2 \alpha_3 \alpha_4 = \exp\left(\frac{\pi}{8} B_C\right) \exp\left(\frac{3\pi}{8} I B_C\right)$$





# Clifford Algebra and the Coxeter Plane – 4D case

rank 4	exponents	W-factorisation
$A_4$	1, 2, 3, 4	$W = \exp\left(\frac{\pi}{5} B_C\right) \exp\left(\frac{2\pi}{5} I B_C\right)$
$B_4$	1, 3, 5, 7	$W = \exp\left(\frac{\pi}{8} B_C\right) \exp\left(\frac{3\pi}{8} I B_C\right)$
$D_4$	1, 3, 3, 5	$W = \exp\left(\frac{\pi}{6} B_C\right) \exp\left(\frac{\pi}{2} I B_C\right)$
$F_4$	1, 5, 7, 11	$W = \exp\left(\frac{\pi}{12} B_C\right) \exp\left(\frac{5\pi}{12} I B_C\right)$
$H_4$	1, 11, 19, 29	$W = \exp\left(\frac{\pi}{30} B_C\right) \exp\left(\frac{11\pi}{30} I B_C\right)$

Actually, in 2, 3 and 4 dimensions it couldn't really be any other way

# Clifford Algebra and the Coxeter Plane – $D_6$


- For  $D_6$  one has exponents  $\boxed{1, 3, 5, 5, 7, 9}$
- Coxeter versor decomposes into orthogonal bits as

$$W = \frac{1}{\sqrt{5}}(e_1 + e_2 + e_3 - e_4 - e_5)e_6 \exp\left(\frac{\pi}{10}B_C\right) \exp\left(\frac{3\pi}{10}B_2\right)$$

- Now **bivector exponentials** correspond to **rotations in orthogonal planes**
- **Vector** factors correspond to **reflections**
- For odd  $n$ , there is always **one such vector factor** in  $D_n$ , and for even  $n$  there are **two**

# Projection and Diagram Foldings

Compare with the 'partial folding' of  $E_8$  to  $H_4$



$$s_{\beta_1} = s_{\alpha_1} s_{\alpha_7}, s_{\beta_2} = s_{\alpha_2} s_{\alpha_6}, s_{\beta_3} = s_{\alpha_3} s_{\alpha_5}, s_{\beta_4} = s_{\alpha_4} s_{\alpha_8} \Rightarrow H_4$$

## Imaginary differences – different imaginaries

So what has been **gained** by this **Clifford view**?

- There are **different** entities that serve as **unit imaginaries**
- They have a **geometric** interpretation as an **eigenplane of the Coxeter element**
- These don't need to **commute** with everything like  $i$  (though they do here – at least anticommute. But that is because we looked for **orthogonal decompositions**)
- But see that in general **naive complexification** can be a dangerous thing to do – **unnecessary**, issues of **commutativity**, **confusing** different imaginaries etc

# Conformal geometry and Clifford algebra

- The conformal group  $C(p, q) \sim SO(p+1, q+1)$
- So can use **versor representation** of conformal transformations in Clifford algebra (**reflections, translations, inversions ...**)
- Treat all of them **multiplicatively** in terms of versors and use **sandwiching**  $Ax\tilde{A}$
- E.g. can generate a whole **root lattice** multiplicatively with **compact reflection part** and **translations**

# Conformal Clifford Algebra

- The **conformal group**  $C(n, p)$  is homomorphic to  $Spin(n+1, p+1)$
- Go to  $e_1, e_2, e_3, e, \bar{e}$ , with  $e_i^2 = 1, e^2 = 1, \bar{e}^2 = -1$
- Define two null vectors  $n \equiv e + \bar{e}, \bar{n} \equiv e - \bar{e}$
- Can embed the 3D vector  $x = x^\mu e_\mu = xe_1 + ye_2 + ze_3$  as a **null vector in 5D** ( $\hat{X} \cdot n = -1$ )

$$F(x) \equiv \hat{X} = \frac{1}{2\lambda^2}(x^2 n + 2\lambda x - \lambda^2 \bar{n})$$

- Essentially **linear** action of  $SO(n+1, p+1)$  in embedding space induces a **non-linear** realisation of the conformal group on the **projective light cone** (Dirac/Hestenes/Lasenby)
- So neat thing is that **conformal transformations** are now done by **rotors** (except inversion which is a reflection) – distances are given by **inner products**

# Operations in Conformal Geometric Algebra

- **Amsterdam protocol:**  $e = e_+$ ,  $\bar{e} = e_-$ ,  $n = n_\infty$  and  $\bar{n} = n_0$ .
- **Reflections**  $y' = -xyx$  since  $e$  and  $\bar{e} \Rightarrow n$  and  $\bar{n}$  are orthogonal to  $x \Rightarrow$  anticommute  $-xnx = n$  and  $-x\bar{n}x = \bar{n}$ :

$$-xF(y)x = F(y') = F(-xyx)$$

- **Rotations**  $y' = Ry\tilde{R}$  from reflections via **Cartan-Dieudonné**



$$RF(y)\tilde{R} = F(y') = F(Ry\tilde{R})$$

- **Translations**  $y' = y + a$  rotor  $T_a = \exp\left(\frac{na}{2\lambda}\right) = 1 + \frac{na}{2\lambda}$

$$T_a F(y) \tilde{T}_a = F(y') = F(y + a)$$

# Proof of Principle

Construction of **root systems** and **quasicrystalline point arrays** carries through, e.g. here for  $H_2$  and a pentagon with translation  $1/\tau$





## Benefits of this approach

- **Conceptual Unification** of Rotations and Translations via rotors
- Construct **root system** from the simple roots as before, and likewise for **quasicrystalline point arrays**
- Increased **numerical stability** (not really an issue here) due to projective representation

# A new set of Bianchi IX Killing Vectors

- Used **Conformal Clifford algebra** setup to treat conformal group  $C(1,3)$  as  $SO(2,4)$
- Stabiliser subgroup of a certain vector gives the **de Sitter** group (Killing vectors)
- Using a certain **projection** broke this down to **two commuting  $SU(2) \times SU(2)$**
- This is a **new set of Bianchi IX Killing vectors** with **nice symmetry properties**

# A unified framework for polyhedral groups

Group	Discrete subgroup	Action Mechanism
$SO(3)$	rotational (chiral)	$x \rightarrow \tilde{R}xR$
$O(3)$	reflection (full/Coxeter)	$x \rightarrow \pm \tilde{A}xA$
$Spin(3)$	binary	$(R_1, R_2) \rightarrow R_1 R_2$
$Pin(3)$	pinor	$(A_1, A_2) \rightarrow A_1 A_2$

- e.g. the **chiral icosahedral** group has 60 elements, encoded in Clifford by 120 rotors, which form the **binary icosahedral** group
- together with the **inversion/pseudoscalar**  $I$  this gives 60 rotations and 60 rotoinversions, i.e. the **full icosahedral** group  $H_3$  in 120 elements (with 240 versors)
- all three are interesting groups, e.g. in **neutrino and flavour physics** for family symmetry model building

## Some Group Theory: chiral, full, binary, pin

- Easy enough to calculate **conjugacy classes** etc of versors in Clifford
- Chiral (**binary**) polyhedral groups have irreps
- tetrahedral (12/24):  $1, 1', 1'', 2_s, 2'_s, 2''_s, 3$
- octahedral (24/48):  $1, 1', 2, 2_s, 2'_s, 3, 3', 4_s$
- icosahedral (60/120):  $1, 2_s, 2'_s, 3, \bar{3}, 4, 4_s, 5, 6_s$
- All binary are **discrete subgroups of  $SU(2)$**  and all thus have a  $2_s$  spinor irrep
- See **McKay correspondence**
- Interesting to look at spinors/binary groups in their own right  
– see **Induction Theorem**

## Some Group Theory: chiral, full, binary, pin

- Full (**Coxeter**) is just **two copies** of this (24/48/120 i.e. same order as binary since both  $\text{Spin}(3)$  and  $O(3)$  are double covers of  $SO(3)$ )
- **Pin group** is just  $1 + I$  of this for  $B_3$  and  $H_3$ , which contain the inversion  $I$
- but **not for  $A_3$ !** (which doesn't – c.f. quaternionic reps)
- Instead  $\text{Pin}(A_3)$  has the same conjugacy classes as  $\text{Spin}(B_3)$

## Conjugacy Classes: Quaternion group $Q$

- Five conjugacy classes:  $\{1\}$ ,  $\{-1\}$ ,  $\{\pm e_1 e_2\}$ ,  $\{\pm e_2 e_3\}$ ,  $\{\pm e_3 e_1\}$
- Different conjugacy classes correspond to different geometric subspaces in the Clifford algebra
- Bit trivial for the quaternion group, but extends to arbitrary dimension

# Conjugacy Classes: Binary octahedral group $2O$

- Eight conjugacy classes:  $\{1\}$ ,  $\{-1\}$ ,  $\underline{6}$ : **bivectors**  $\{\pm e_1 e_2, \pm e_2 e_3, \pm e_3 e_1\}$ ;  $\underline{6}'$ : **bivector exponentials**  $\exp \underline{6}$ ;  $\underline{6}''$ :  $\exp -\underline{6}$ ;  $\underline{8}$ : **spinors**  $\{1 \pm e_1 e_2 \pm e_2 e_3 \pm e_3 e_1, \dots\}$ ;  $\underline{8}'$ :  $-\underline{8}$ ;  $\underline{12}$ : **bivectors**  $\{e_1(e_2 + e_3), \dots\}$ ;
- Turns out most of these are the **same as for  $\text{Pin}(A_3)$** , and the remaining ones can be mapped to each other
- Though in  $\text{Pin}(A_3)$  also have **odd grade** elements, so some of the conjugacy classes are vector+trivector etc, i.e. different geometric interpretation

# Character tables: Quaternion group $Q$ (from $A_1^3 \rightarrow A_1^4$ )

$Q$	1	-1	$\pm i$	$\pm j$	$\pm k$
1	1	1	1	1	1
$1'$	1	1	1	-1	-1
$1''$	1	1	-1	1	-1
$1'''$	1	1	-1	-1	1
2	2	-2	0	0	0
4	4	-4	0	0	0

Latter is of **quaternionic type** – somehow seen as particularly noteworthy



# Character tables: binary octahedral group $2O$ (from $B_3 \rightarrow F_4$ )

$2O$	1	1	6	8	8	6	6	12
1	1	1	1	1	1	1	1	1
$1'$	1	1	1	1	1	-1	-1	-1
2	2	2	2	-1	-1	0	0	0
3	3	3	-1	0	0	1	1	-1
$3'$	3	3	-1	0	0	-1	-1	1
4	4	-4	0	2	-2	$2\sqrt{2}$	$-2\sqrt{2}$	0
$4'$	4	-4	0	2	-2	$-2\sqrt{2}$	$2\sqrt{2}$	0
8	8	-8	0	-2	2	0	0	0

Again some of quaternionic type

# Representations

- This **Clifford multivector construction** of the polyhedral groups is a **faithful realisation/representation**, i.e. is essentially the same as the abstract group
- But can define several different **representations** from these versor groups (may or may not be **irreducible** ones)
- Representations: matrices  $D(R)$  such that

$$D(R_1 R_2) = D(R_1) D(R_2)$$

# Representations

- Representations: matrices  $D(R)$  such that

$$D(R_1 R_2) = D(R_1) D(R_2)$$

- **Trivial** representation:  $D(R) = R1\tilde{R} = 1$
- **Rotation** representation: for nD vector  $x = \sum a_i e_i$ :

$$D(R)\underline{x} = R x \tilde{R} \quad \text{usual } SO(n) \text{ } n \times n\text{-matrix}$$

- Full representation: for nD vector  $x = \sum a_i e_i$ :

$$D(A)\underline{x} = A x \tilde{A}$$

usual  $O(n)$   $n \times n$ -matrix

- **Spinor** representation: for nD spinor  $y$  ( $2^{n-1}$  components):

$$D(R)\underline{y} = R y \quad \text{a } 2^{n-1} \times 2^{n-1}\text{-matrix}$$

- **Versor** representation: for nD versor  $z$  ( $2^n$  components):

$$D(A)\underline{z} = A z \quad \text{a } 2^n \times 2^n\text{-matrix}$$

# Character tables and Clifford reps: quaternion group $Q$

The **spinor** representation  $D(R)\underline{y} = R\underline{y}$  of the quaternion group  $Q$  gives the representation of **quaternionic type**. (The **trace** of  $D(R)$  is the **character**.)

Again just seen to be a consequence of the **accidental isomorphism** between 3D spinors and quaternions.

$Q$	1	-1	$\pm i$	$\pm j$	$\pm k$
1	1	1	1	1	1
1'	1	1	1	-1	-1
1''	1	1	-1	1	-1
1'''	1	1	-1	-1	1
2	2	-2	0	0	0
4	4	-4	0	0	0

# Character tables and Clifford reps: binary octahedral group $2O$

The **spinor** representation  $\boxed{D(R)\underline{y} = Ry}$  of the quaternion group  $2O$  gives the irrep of **quaternionic type**.

The **rotation** representation  $\boxed{D(R)\underline{x} = Rx\tilde{R}}$  gives **3 irrep**.

$2O$	1	1	6	8	8	6	6	12
1	1	1	1	1	1	1	1	1
$1'$	1	1	1	1	1	-1	-1	-1
2	2	2	2	-1	-1	0	0	0
3	3	3	-1	0	0	1	1	-1
$3'$	3	3	-1	0	0	-1	-1	1
4	4	-4	0	2	-2	$2\sqrt{2}$	$-2\sqrt{2}$	0
$4'$	4	-4	0	2	-2	$-2\sqrt{2}$	$2\sqrt{2}$	0
8	8	-8	0	-2	2	0	0	0

# Clifford: groups and representations summary

- Clifford algebra provides a **unified framework** for chiral/full/binary/pin **polyhedral** groups all in the **same** representation/realisation space/**algebra** (c.f. usual  $SO(3)$  vs  $SU(2)$  representations)
- Structure of the algebra  $\Rightarrow$  different **conjugacy classes** are **different kinds of objects in the algebra** and are kept separate
- Several **representations** follow, in particular have **geometric insight** into **complex and quaternionic** representations

## Clifford summary

- Interesting **induction theorem** linking geometry of 3D and 4D
- Geometric **complex structures** and **non-trivial commutativity** properties
- Simple versor representation of **orthogonal transformations**
- **Conformal** geometry
- Some interesting new results on **group and representation theory**
- **Lie algebras** can be constructed in Clifford algebra as **bivector algebras**, and **Lie groups** as **spin groups** (work with Phoenix)

## 1 Introduction

- Coxeter groups and root systems
- Clifford algebras

## 2 Coxeter and Clifford

- The Induction Theorem – from 3D to 4D
- The Coxeter Plane
- Conformal Geometry
- Some Group Theory

## 3 Moonshine and Outlook



# Monstrous Moonshine

- **Mysterious connection** between two very different areas of Mathematics
- **Modular forms** (functions that live on a torus with complex structure  $\tau$ ): Fourier expansion **coefficients** wrt ( $q = e^{2\pi i\tau}$ )
- **Finite simple groups**: **dimensions** of irreducible representations
- **Monstrous Moonshine**: The largest sporadic group, the **Monster  $M$**  ( $\{1, 196883, 21296876, \dots\}$ ) and the Klein  **$j(\tau)$**  modular function
- $j(\tau) = q^{-1} + 744 + 196884q + 21493760q^2 + \dots$
- $196884 = 196883 + 1, 21493760 = 21296876 + 196883 + 1, \dots$

# Mathieu Moonshine

- Similar Moonshine phenomenon
- Modular form: elliptic genus of an  $\mathcal{N} = 4$  SCFT compactified on a K3-surface
- Finite simple group: Mathieu  $M_{24}$   
( $\{45, 231, 770, 2277, 5796 \dots\}$ )
- Elliptic genus is

$$E_{K3}(\tau, z) = -2Ch(0; \tau, z) + 20Ch(1/2; \tau, z) + e(q)Ch(\tau, z)$$

where all the coefficients in the  $q$ -series

$$e(q) = 90q + 462q^2 + 1540q^3 + 4554q^4 + 11592q^5 + \dots$$

are twice the dimension of some  $M_{24}$  irrep

# Clifford and Moonshine

- Looking at **Wess-Zumino-Witten models**, i.e. strings propagating on a Lie group manifold
- Condition of **extended supersymmetry** ultimately hinges on **classification** of Clifford algebras – deep connection
- Connections with **binary polyhedral groups**, **Monstrous Moonshine**, **McKay correspondence**, **lattices**, **affine extensions**, **Lie groups/algebras** etc
- **Elliptic genus** is constant on (connected components of) the **moduli space**: better understanding as a **topological** feature as the **index of a Dirac operator**?

Thank you!

## Back to the roots

- Unifying principles of **geometry** and **symmetry**
- **Discrete** groups (finite simple groups, polyhedral groups) and **continuous** groups (string compactifications, non-linear  $\sigma$ -models)
- **Exceptional phenomena**:  $E_8$ ,  $H_4/F_4/D_4$  (spinorial), McKay correspondence, Monster  $M$ , Mathieu  $M_{24}$  ...
- **Applications**: from **mundane** (viruses, fullerenes, quasicrystals) to **exotic** (HEP, Moonshine) – **same mathematical tools**: Coxeter, Clifford, affine extensions etc


# Mathematical Aspects: Better understanding of geometrical structures

- Affine extensions and translations – lattices, quasicrystals and projections
- Coxeter groups and Kac-Moody theory
- Euclidean, spherical, hyperbolic and conformal geometry
- Clifford and spinorial geometry
- Group theory, Lie groups and algebras (with Phoenix)
- Mathieu Moonshine, McKay correspondence
- Implications for the real world

# HEP

- Gravitational and cosmological singularities – hidden symmetries
- New uses for non-crystallographic groups
- Topological defects
- Integrable systems
- Family symmetries (flavour/neutrino physics)
- 4D groups:  $A_4$ ,  $B_4$ ,  $D_4 - F_4$ ,  $H_4$ ?
- Spinor geometry
- $E_8$  features prominently – relation with  $H_4$

# Projection and Diagram Foldings



$$s_{\beta_1} = s_{\alpha_1} s_{\alpha_7}, s_{\beta_2} = s_{\alpha_2} s_{\alpha_6}, s_{\beta_3} = s_{\alpha_3} s_{\alpha_5}, s_{\beta_4} = s_{\alpha_4} s_{\alpha_8} \Rightarrow H_4$$

$E_8$  has two  $H_4$ -invariant subspaces – blockdiagonal form

$D_6$  has two  $H_3$ -invariant subspaces

$A_4$  has two  $H_2$ -invariant subspaces






## 4D geometry is surprisingly important for HEP

- 4D root systems are **surprisingly relevant to HEP**
- $A_4$  is  $SU(5)$  and comes up in **Grand Unification**
- $D_4$  is  $SO(8)$  and is the little group of **String theory**
- In particular, its **triality symmetry** is crucial for showing the equivalence of RNS and GS strings
- $B_4$  is  $SO(9)$  and is the little group of **M-Theory**
- $F_4$  is the **largest crystallographic** symmetry in 4D and  $H_4$  is the **largest non-crystallographic** group
- The above are **subgroups** of the latter two
- **Spinorial nature** of the root systems could have **surprising consequences for HEP**


# Motivation: Viruses

- Geometry of **polyhedra** described by **Coxeter** groups
- Viruses have to be 'economical' with their **genes**
- Encode **structure** modulo **symmetry**
- **Largest discrete symmetry of space** is the **icosahedral** group
- Many other 'maximally symmetric' objects in nature are also icosahedral: **Fullerenes & Quasicrystals**
- But: viruses are not just polyhedral – they have **radial structure**. **Affine extensions** give **translations**



## Extend icosahedral group with distinguished translations




- Radial layers are **simultaneously constrained** by affine symmetry
- Works very well in practice: **finite library of blueprints**
- **Select** blueprint from the **outer shape** (capsid)
- Can **predict inner structure** (nucleic acid distribution) of the virus from the point array



**Affine extensions** of the icosahedral group (giving translations) and their **classification**.


# Use in Mathematical Virology

- Suffice to say **point arrays work very exceedingly well** in practice. Two papers on the mathematical (Coxeter) aspects.
- **Implemented computational problem in Clifford** – some **very interesting mathematics** comes out as well (see later).




## Extension to fullerenes: carbon onions

- Extend idea of affine symmetry to other icosahedral objects in nature: football-shaped **fullerenes**
- Recover different shells with icosahedral symmetry from affine approach: **carbon onions** ( $C_{60} - C_{240} - C_{540}$ )




## Extension to fullerenes: carbon onions

- Extend idea of affine symmetry to other icosahedral objects in nature: football-shaped **fullerenes**
- Recover different shells with icosahedral symmetry from affine approach: **carbon onions** ( $C_{80} - C_{180} - C_{320}$ )



## Applied areas (EPSRC proposal)

- **Viruses:** Extend to large viruses; interesting results on **higher-order translations**
- **Proteins:** Extend affine symmetry to  **$2D$**  and apply to (**chiral**) proteins
- **Fullerenes:** Extend to larger fullerenes, in particular **chiral carbon onions**
- **Packings:** Novel analytical and numerical approaches to packings of polyhedral solids; with Colapinto (Santa Barbara), Twarock (York) and Thorpe (Phoenix)



# References

- Novel Kac-Moody-type affine extensions of non-crystallographic Coxeter groups with Twarock/Bøhm  
J. Phys. A: Math. Theor. 45 285202 (2012)
- Affine extensions of non-crystallographic Coxeter groups induced by projection with Twarock/Bøhm  
Journal of Mathematical Physics 54 093508 (2013), [Cover article September](#)
- Viruses and Fullerenes – Symmetry as a Common Thread?  
with Twarock/Wardman/Keef March Cover Acta Crystallographica A (2014)